

**WestminsterResearch**

<http://www.westminster.ac.uk/westminsterresearch>

**Investigating mitochondrial DNA relationships in Neolithic  
Western Europe through serial coalescent simulations**

**Rivollat, M., Rottier, S., Couture, C., Pemonge, M.H., Mendisco, F.,  
Thomas, M.G., Deguilloux, M.F. and Gerbault, P.**

This is an author's accepted manuscript of an article published in the European Journal of Human Genetics, 28 December 2016. The final definitive version is available online at:

<https://dx.doi.org/10.1038/ejhg.2016.180>

---

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

---

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch: (<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail [repository@westminster.ac.uk](mailto:repository@westminster.ac.uk)

1 SHORT REPORT

2  
3 Title:

4 Investigating mitochondrial DNA relationships in Neolithic Western Europe through  
5 serial coalescent simulations

6  
7 Running title:

8 Genetic relationships in Neolithic western Europe  
9

10  
11 Authors:

12 Maïté Rivollat<sup>1</sup>, Stéphane Rottier<sup>1</sup>, Christine Couture<sup>1</sup>, Marie-Hélène Pemonge<sup>1</sup>, Fanny  
13 Mendisco<sup>1</sup>, Mark G. Thomas<sup>2</sup>, Marie-France Deguilloux<sup>1</sup>, Pascale Gerbault<sup>2,3,4</sup>  
14

15 Affiliations:

16 <sup>1</sup> De la Préhistoire à l'Actuel, Culture, Environnement, Anthropologie – UMR 5199,  
17 CNRS, Université de Bordeaux, Bordeaux, France

18 <sup>2</sup> UCL Research Department of Genetics, Evolution and Environment, Darwin building,  
19 Gower Street, London WC1E 6BT, UK

20 <sup>3</sup> UCL Department of Anthropology, 14 Taviton Street, London WC1H 0BW, UK

21 <sup>4</sup> Department of Life Sciences, University of Westminster, 115 New Cavendish Street,  
22 London W1W 6UW, UK  
23

24 Corresponding authors:

25 Maïté Rivollat (email : maite.rivollat@u-bordeaux.fr / maite87@live.fr; Tel: +33 (0)5  
26 40 00 25 48; Fax: +33 (0)5 40 00 25 45); Pascale Gerbault (email:  
27 p.gerbault@ucl.ac.uk; Tel: +44 (0) 207 679 4397; Fax: +44 (0) 207 679 7193)  
28

29 Conflict of interest:

30 The authors have nothing to disclose, no conflict of interest.  
31  
32  
33

Abstract:

Recent ancient DNA studies on European Neolithic human populations have provided persuasive evidence of a major migration of farmers originating from the Aegean, accompanied by sporadic hunter-gatherer admixture into early Neolithic populations, but increasing towards the Late Neolithic. In this context, ancient mitochondrial DNA (mtDNA) data collected from the Neolithic necropolis of Gurgy (Paris Basin, France), the largest mtDNA sample obtained from a single archaeological site for the Early/Middle Neolithic period, indicate little differentiation from farmers associated to both the Danubian and Mediterranean Neolithic migration routes, as well as from western European hunter-gatherers. To test whether this pattern of differentiation could arise in a single unstructured population by genetic drift alone, we used serial coalescent simulations. We explore female effective population size parameter combinations at the time of the colonization of Europe 45 000 years ago and the most recent of the Neolithic samples analyzed in this study 5 900 years ago, and identify conditions under which population panmixia between hunter-gatherers/Early-Middle Neolithic farmers and Gurgy cannot be rejected. In relation to other studies on the current debate of the origins of Europeans, these results suggest increasing hunter-gatherer admixture into farmers' group migrating farther west in Europe.

Key words: genetic drift, European Neolithic, serial coalescent, ancient DNA, mtDNA

Introduction:

The introduction of farming into Europe around 8 600 years ago led to fundamental changes in subsistence strategy and social organization, and left signatures of

population turnover<sup>1-4</sup>. It is widely believed that farming spread into Europe from the Aegean along both Mediterranean and Danubian routes<sup>3,5</sup>. Recent archaeological<sup>6</sup> and palaeogenetic evidence<sup>1,2,4,5,7</sup> indicate a crucial role for migration, with only sporadic hunter-gatherer (HG) admixture into early Neolithic populations, but increasing towards the Late Neolithic<sup>1,7,8</sup>. However, these local inferences still permit spatiotemporal heterogeneity in HG admixture during the Neolithic in continental Europe.

In this context, the mtDNA diversity of the Gurgy "Les Noisats" site, located south of the Paris Basin and dated from 7 000 to 6 000 years ago, is striking since descriptive analyses<sup>9</sup> indicated affinities not only with early farmers associated with both the Danubian and Mediterranean migration routes but also with European HG. Notably, a relatively lower differentiation between Gurgy and European HG ( $F_{ST}=0.08$ ) was observed when compared to other published levels of differentiation between Early Neolithic farmers and HG (e.g.  $F_{ST}=0.0923$  (ref. 10);  $F_{ST}=0.163$  (ref. 2)). This suggests complex admixture pattern between HG and farmer groups to shape Gurgy mtDNA diversity.

Previous mtDNA studies<sup>2,10,11</sup> have used serial coalescent simulations to test for genetic continuity between HG, Neolithic farmers and extant DNA samples from the same geographic region, and regularly concluded in genetic discontinuity between groups.

We used a similar approach to address if the observed level of mtDNA differentiation between European HG, Neolithic farmer and Gurgy groups could be obtained under a panmictic population model with various combinations of effective population sizes.

Our approach differs in three major aspects from previous studies<sup>2,10,11</sup>: first, we grouped the ancient mtDNA sequences according to subsistence strategy (HG or Neolithic farmers) and Neolithic context (Mediterranean/South-, Danubian/Central- or

Gurgy- farmers). Some of the sample groups are consequently contemporaneous and can represent various regions. Second, we did not include modern population sample into the comparison. Third, we extended the effective population size ranges used previously<sup>2,11</sup> towards the lower bound to explore further demographic scenarios.

## MATERIAL AND METHODS

We compiled 282 available ancient mtDNA HVR-I sequences (NC\_012920.1:m.16024\_16380; Table S1). Following Rivollat et al. 2015, ancient mtDNA data were partitioned into 4 sample groups: (i) Gurgy Les Noisats necropolis (hereafter referred to as “Gurgy”,  $n=39$  sequences), (ii) Neolithic farmers from south Europe (group “South-F”,  $n=56$ , partitioned into 4 chronological sub-groups), (iii) Neolithic farmers from central Europe (“Central-F”,  $n=147$ , 5 sub-groups), and (iv) hunter-gatherers (“HG”,  $n=40$ , 16 sub-groups). Chronological sub-groups were defined according to both shared geographic location and median calibrated C14 dates (see Figure 1 and Figure 2). As a test statistic that measures the level of population differentiation, we calculated six pairwise  $F_{ST}$  between the four groups (Figure 3) with ARLSUMSTAT version 3.5.1.2 (ref. 12).

Following previous studies<sup>2,11</sup> we performed serial coalescent simulations under a single panmictic population model with two demographic events: an initial colonization of Europe 45 000 years ago of female effective population size  $N_{UP}$ , followed by exponential growth or decline to the Neolithic transition in Western Europe 5 900 years cal. BP of female effective population size  $N_N$ . Prior to  $N_{UP}$  we assume an ancestral female effective population size  $N_A$  of 5 000, derived from the commonly used long-term effective human population size of 10 000 individuals outside Africa<sup>13</sup> and

assuming a 1:1 female to male ratio. We explored 50 values for  $N_{UP}$  ranging from 1 to 5  
 000 and 50 values for  $N_N$  ranging from 10 to 100 000 (Table S2). We generated 50 000  
 mitochondrial genealogies of ancient HG and farmer sequences using fastsimcoal  
 version 2.5.1 (ref. 14) under each of the 2 500  $N_{UP}$  -  $N_N$  combinations (Table S2). We  
 used a fixed mutation rate of  $5 \times 10^{-6}$ /bp/generation (ref. 15), assuming a 25 years  
 generation time. These simulated genealogies were used to compute expected pairwise  
 $F_{ST}$  values for the six sample comparisons (Figure 2). We recorded the proportion of  
 simulated  $F_{ST}$  values that were greater than those observed per  $F_{ST}$  and parameter  
 combination (Figure 3).

We also tested if the six observed pairwise  $F_{ST}$  values as well as eight within sample  
 group statistic values (number of segregating sites and of pairwise differences) could be  
 recovered from simulations under this simple model by performing an approximate  
 Bayesian computation (ABC) -related approach<sup>16</sup> (see details in SI). We used the  
 rejection algorithm of the ‘abc’ package<sup>17</sup> available in R to retain the parameter  
 combinations that generated simulated pairwise  $F_{ST}$  the closest to the 6 observed values.  
 Even though we provide some effective population size estimates, we caution against  
 over-interpretation since there is likely insufficient information in the data to make  
 precise estimates.

## Results and Discussion

Analyses indicate that for the six pairwise population group comparisons, some  $N_{UP}$  -  
 $N_N$  combinations can result in simulated differentiation greater than the one observed  
 (grey area on Figure 3). Notably, results show that we cannot reject the possibility that

European HG, South-F, Central-F and Gurgy were sampled from a single panmictic population. Whereas these results may appear to contrast with previous studies that have used serial coalescent simulations to address local mtDNA population continuity between diachronic HG and farmers samples<sup>2,11</sup>, we highlight that our analyses do not address ‘population continuity’ as defined in these studies. The grouping of diachronic samples may artificially reduce the level of differentiation that would be observed in case of significant mtDNA population structure. This grouping none-the-less allows us to investigate the genetic relationships between set of lineage samples associated with specific archaeological Neolithic contexts.

We confirmed that our panmictic population model generated simulated between and within population group diversity values close to the observed using an ABC-rejection algorithm (see SI and Figure S1). The 95% credible intervals estimated from the retained simulations are [5 – 3500]  $N_{UP}$  females and [200 – 7750]  $N_N$  females. These estimates concur with the observation that the parameter space for which a panmictic population model may hold is rather narrow (Figure 3). Most  $N_N$  values tested and compatible with the level of mtDNA differentiation observed are relatively low (10 to 200 females for the South-F and Central-F comparison, Figure 3). Noteworthy, some  $N_{UP}$  -  $N_N$  combinations imply a population decline that clearly contrasts with previous studies based on modern DNA data which have inferred female effective population size growth in Europe during the Holocene<sup>18</sup>. However, we were not constrained to simulate population expansion, since we did not consider modern DNA data in our analyses. Moreover, a Holocene population decline in Europe corroborates recent Y chromosome data<sup>18</sup> and various archaeological evidence support demographic fluctuation of Neolithic populations<sup>19,20</sup>.

Our results indicate that a simple panmictic population model can account for the mtDNA differentiation observed between European HG and Early/Middle Neolithic farmers; a larger proportion of the HG - Gurgy explored parameter space failed to reject panmixia. This result suggests increasing HG admixture into farmers' group migrating farther west in Europe. Similarly, we note that a larger proportion of the explored parameter space fails to reject panmixia when comparing Gurgy and South-F than when comparing Gurgy and Central-F. Thus, our results seem to support Gurgy as the most ancient Neolithic sample studied so far with appreciable admixture between pre-Neolithic HG and Early/Middle Neolithic farmers from both streams of Neolithization in Europe (with a suspected higher participation of Mediterranean farmers).

As with any model, the one we test here has a few assumptions that may not hold, e.g.  $N_A$  of female to male ratio of 1 (ref. 18) and no population structure in any of the four groups<sup>5</sup>. Moreover, the panmictic population model proposed would need to be compared against alternatives (e.g. ref. 11). Such a simple panmictic population model nevertheless lays the ground for building more complex ones<sup>17</sup>. Notably, a serial coalescent approach coupled with ABC would allow estimation of the possible contribution of each of the three population groups (HG, Mediterranean and Central Europe farmers) in shaping Gurgy mitochondrial diversity.

#### Acknowledgements:

This work was supported by a ministerial grant from the Research National Agency as a program of prospects investments (ANR-10-LABX-52, DHP project; dir: SR; Université Bordeaux 1, LaScArBx-ANR; 2012-14; and ANR-10-IDEX-03-02 to MR



for work in UCL) and a Leverhulme Programme grant to A.M. Migliano (UCL Anthropology) and M.G. Thomas (RP2011-R-045). The authors acknowledge the use of the UCL Legion High Performance Computing Facility (Legion@UCL), and associated support services, in the completion of this work.

Supplementary information is available at *European Journal of Human Genetics'* website.

## References

- 1 Haak, W., Lazaridis, I., Patterson, N. *et al.* Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature* 2015; **552**, 207-211.
- 2 Bramanti, B., Thomas, M. G., Haak, W. *et al.* Genetic discontinuity between local hunter-gatherers and central Europe's first farmers. *Science* 2009; **326**, 137-140, doi:10.1126/science.1176869.
- 3 Childe, V. G. *The Dawn of European Civilization*. Vol. 674 Kegan Paul, 1925; 328.
- 4 Lazaridis, I., Patterson, N., Mitnik, A. *et al.* Ancient human genomes suggest three ancestral populations for present-day Europeans. *Nature* 2014; **513**, 409-413.

200 5 Hofmanová, Z., Kreutzer, S., Hellenthal, G. *et al.* Early farmers from across  
201 Europe directly descended from Neolithic Aegeans. *Proceedings of the*  
202 *National Academy of Sciences* in press.

203 6 Bocquet-Appel, J.-P., Naji, S., Vander Linden, M. & Kozłowski, J. K. Detection  
204 of diffusion and contact zones of early farming in Europe from the space-  
205 time distribution of 14C dates. *Journal of Archaeological Science* 2009; **36**,  
206 807-820.

207 7 Brandt, G., Haak, W., Adler, C. J. *et al.* Ancient DNA Reveals Key Stages in the  
208 Formation of Central European Mitochondrial Genetic Diversity. *Science*  
209 2013; **342**, 257-261, doi:10.1126/science.1241844.

210 8 Bollongino, R., Nehlich, O., Richards, M. P. *et al.* 2000 years of parallel  
211 societies in Stone Age Central Europe. *Science* 2013; **342**, 479-481,  
212 doi:10.1126/science.1245049.

213 9 Rivollat, M., Mendisco, F., Pemonge, M.-H. *et al.* When the Waves of  
214 European Neolithization Met: First Paleogenetic Evidence from Early  
215 Farmers in the Southern Paris Basin. *PLoS One* 2015; **10**, e0125521,  
216 doi:10.1371/journal.pone.0125521.

217 10 Haak, W., Balanovsky, O., Sanchez, J. J. *et al.* Ancient DNA from European  
218 early neolithic farmers reveals their near eastern affinities. *PLoS biology*  
219 2010; **8**, e1000536, doi:10.1371/journal.pbio.1000536.

220 11 Rasteiro, R. & Chikhi, L. Female and male perspectives on the neolithic  
221 transition in Europe: clues from ancient and modern genetic data. *PLoS One*  
222 2013; **8**, e60944, doi:10.1371/journal.pone.0060944.

223 12 Excoffier, L. & Lischer, H. E. Arlequin suite ver 3.5: a new series of programs  
224 to perform population genetics analyses under Linux and Windows.  
225 *Molecular ecology resources* 2010; **10**, 564-567.

226 13 Consortium, T. G. P. A map of human genome variation from population-  
227 scale sequencing. *Nature* 2010; **467**, 1061-1073.

228 14 Excoffier, L. & Foll, M. Fastsimcoal: a continuous-time coalescent simulator  
229 of genomic diversity under arbitrarily complex evolutionary scenarios.  
230 *Bioinformatics* 2011; **27**, 1332-1334.

231 15 Soares, P., Ermini, L., Thomson, N. *et al.* Correcting for purifying selection:  
232 an improved human mitochondrial molecular clock. *The American Journal of*  
233 *Human Genetics* 2009; **84**, 740-759.

234 16 Beaumont, M. A., Zhang, W. & Balding, D. J. Approximate Bayesian  
235 computation in population genetics. *Genetics* 2002; **162**, 2025-2035.

236 17 Csilléry, K., Blum, M. G., Gaggiotti, O. E. & François, O. Approximate Bayesian  
237 computation (ABC) in practice. *Trends in Ecology & Evolution* 2010; **25**, 410-  
238 418.

239 18 Karmin, M., Saag, L., Vicente, M. *et al.* A recent bottleneck of Y chromosome  
240 diversity coincides with a global change in culture. *Genome Research* 2015;  
241 **25**, 459-466.

242 19 Shennan, S., Downey, S. S., Timpson, A. *et al.* Regional population collapse  
243 followed initial agriculture booms in mid-Holocene Europe. *Nature*  
244 *communications* 2013; **4**, 2486.

245 20 Wirtz, K. W. & Lemmen, C. A global dynamic model for the Neolithic  
246 transition. *Climatic Change* 2003; **59**, 333-367.

247

248


‘Titles and legends to figures’:

Figure 1: Geographic locations of the sites sampled for ancient mtDNA in the current analysis. 1 refers to the most recent sample group in the gene genealogy and 26 to the most ancient. Groups were categorized based on common median C14 dates, archaeological context and geographic location; one group can consequently be sampled from more than one geographic location; for those groups with more than one location, only the location of the largest sample size group is displayed on the map (details in Figure 2 and Table S1).

Figure 2: Demographic model simulated with the serial coalescent. Time is the median calibrated C14 years before present (cal BP) backward in time from ‘t0’ and expressed in generations. ‘t0’ refers to 5 900 years cal. BP, the median C14 date of the youngest ancient mtDNA sample. Groups are numbered backward in time from the most recent to the most ancient. The dashed red cylinder shows constant population size between  $N_{UP}$  and  $N_N$ , but the simulated population can undergo expansion or decline depending on the combinations of these parameter values.

Figure 3: Probability of obtaining simulated  $F_{ST}$  value greater than that observed for the six pairwise population groups compared (see text for details). Corresponding observed pairwise  $F_{ST}$  are shown in the top left corner of each grid. The 50x50 grids show values of assumed effective population size  $N_N$  on the x-axis and values of parameterized  $N_{UP}$  on the y-axis (note that 25 values are shown on each axis for clarity, see Table S2). The top right area delimited by vertical and horizontal black lines outline  $N_N$  and  $N_{UP}$  ranges, respectively, used in comparable studies<sup>2,11</sup>. Grey shows proportions of observed  $F_{ST}$  greater than observed (proportion > 0.05), for which panmixia cannot be rejected. Color-scale represents

significance level from blue (proportion lower or equal to 0.05) to red (proportion close to 0). Proportions were obtained over 50 000 simulated pairwise  $F_{ST}$  per combination of  $N_N$  and  $N_{UP}$  value.



Population name (total n)	group	NB	Sample Size (n)	Time (gen back)	Median date (cal. BP)
Gurgy	5	39	24	6500	
South_F (56)	1	3	0	5900	
	6	36	32	6700	
	7	7	42	6950	
	10	10	54	7250	
Central_F (147)	2	32	4	6000	
	4	17	20	6400	
	8	88	44	7000	
	9	7	52	7190	
	12	3	71	7685	
HG (40)	3	3	14	6250	
	11	1	64	7500	
	13	2	76	7800	
	14	11	86	8050	
	15	1	96	8300	
	16	2	96	8300	
	17	2	110	8650	
	18	1	116	8800	
	19	2	144	9500	
	20	2	312	13700	
	21	1	324	14000	
	22	3	344	14500	
	23	1	378	15350	
	24	4	764	25000	
	25	3	1010	31155	
	26	1	1343	39475	

